

ANTIMICROBIAL PEPTIDES: A PROMISING FUTURE ALTERNATIVE TO ANTIBIOTICS IN AQUACULTURE

ANUTOSH PARIA, VINAY T.N., SANJAY K. GUPTA,
TANMOY GON CHOUHURY AND BIPLAB SARKAR

Intensification in aquaculture systems has brought into question the overall well-being of fish, human and the environment. The need to produce more in the culture system imposes unintentional consequences to the fish which weakens the immune system of fish and finally leads to disease outbreak. Furthermore, introduction of non-native fish species also introduces many transboundary aquatic animal diseases. Up to 40 percent of shrimp aquaculture production is lost annually, worth more than \$3 billion, mainly due to viral pathogens (Stentiford *et al.* 2012). Disease losses in aquaculture can account for about 30 percent of total operating costs (Lee and O'Bryen 2003).

To efficiently control disease outbreaks in aquaculture systems, a “prevention is better than treatment” approach based on the principle of prophylaxis is the method of choice. As examples, immunostimulants, probiotics, prebiotics and vaccines are being used for disease prevention. However, especially in intensive culture environments, mass mortality occurs even after taking precautions. To prevent the stock being wiped out after a disease outbreak, use of chemotherapeutics such as antibiotics, antiviral drugs and other chemical therapeutants like formaldehyde, sodium chloride, potassium permanganate are used.

The gross global investment in aquatic animal health products is \$274.4 million for antibiotics and \$29.4 million for antiparasitics (The Fish Site 2010). Apart from these drugs, next-generation antimicrobials and antimicrobial peptides with broad-spectrum activity are in focus as alternate therapeutic agents in aquaculture.

ANTIMICROBIAL RESISTANCE AND NEED FOR NOVEL ANTIMICROBIAL AGENTS IN AQUACULTURE

Antimicrobial drugs have been the most common choice for treating many bacteria-borne diseases in people and livestock, including aquaculture, for over 50 years with tremendous benefits. But inappropriate administration of different classes of antimicrobials in aquaculture provides a selective pressure, creating a reservoir of multiple antimicrobial-resistant bacteria in cultured fish and shrimp and in the culture environment. Antimicrobial resistance is rising alarmingly. In aquaculture, antibiotic




FIGURE 1. The lack of introduction of potent new antibiotics in the latter half of last century. (Adapted from Fischbach and Walsh 2009.)

resistance has been reported in pathogenic bacteria of fish such as *Aeromonas salmonicida*, *A. hydrophila*, *Vibrio anguillarum*, *Pseudomonas fluorescens*, *Pasteurella piscicida*, *Edwardsiella tarda* (Aoki 1988) and *Yersinia ruckeri* (DeGrandis and Stevenson 1985). The presence of multiple antibiotic-resistance bacteria in aquaculture

food products has become a serious human health threat because there is a potential that genes responsible for drug resistance may be transferred to bacteria in humans (Kathleen *et al.* 2016).

Although the number of multi-drug resistant bacteria is increasing, the invention of new antibiotics has been very low. From 1960s to early 2010, only four new classes of antibiotics were introduced and none have made any substantial impact (Fig. 1). The global antibiotic market is still largely dominated by antibiotics discovered before the 1960s and the few antibiotics produced since then were synthetic derivatives of existing core structures (Fischbach and Walsh 2009). In this regard, developing and finding new compounds with broad-spectrum antimicrobial activity is a pressing need. This scenario even worse in the case of aquaculture because only one antibiotic (oxytetracycline hydrochloride) is approved by the US Food and Drug Administration for use against different bacterial diseases. Hence, it is the utmost priority to find an alternative antimicrobial agent to reduce production losses from disease epizootics in aquaculture systems.

ANTIMICROBIAL PEPTIDES AS NOVEL ANTIMICROBIAL AGENT

Antimicrobial peptides (AMPs) are a group of highly conserved oligopeptides with five to over 100 amino acids. They are pivotal humoral components of innate immunity and have been extensively studied in invertebrates and vertebrates including fish and exhibit broad-spectrum antimicrobial activity *in vitro* and *in vivo*. They are cationic molecules that can selectively attack negatively-charged bacterial cell membranes and brings about destruction of the organism by pore formation or destabilization of membrane equilibrium or by penetration into the cell and

(CONTINUED ON PAGE 68)

thereby the impairment of cellular machinery (Reddy *et al.* 2004) (Fig. 2). Because of the wide activity against different microbial agents, these molecules have made their mark as an emerging class of natural antibiotics. Apart from having an antimicrobial role against many microorganisms, the salient features of AMPs are their broad-spectrum antimicrobial activity towards multi-drug resistant microbial isolates and very minimal chance to allow development resistance due to rapid lysis of cell membranes (Lai and Gallo 2009) (Fig. 2). Antimicrobial peptides (AMPs) have been isolated and characterized as a novel and useful alternative to antibiotics (Mookherjee and Hancock 2007). Even disease-resistant transgenic fish stock was developed to target selected AMPs (Buchanan *et al.* 2001, Sarmasik *et al.* 2002).

FISH ANTIMICROBIAL PEPTIDES

Most fish AMPs are potent antibiotics with a broad spectrum of activity at micromole concentrations (Rajanbabu and Chen 2011). Although the first AMP found in fish was purified in 1995 (Pilet *et al.* 1995), it took almost a decade to learn about its role as a major component of the innate immune response.

Piscidins are the family of linear, amphipathic AMPs. The first member of this family was identified from skin mucous secretions of the winter flounder *Pleuronectes americanus* and called pleurocidin (Cole *et al.* 1997). Other piscidins homologous to pleurocidins were identified in other teleosts (named asmisgurin, moronecinin, epinecidin and dicentracin). Generally, piscidins are found in gill, skin and intestine of finfish. They have potent antimicrobial activity against a variety of microorganisms. These are specifically effective against Gram-positive and -negative bacteria, with the best antimicrobial activity observed in the case of *Streptococcus*, *Pseudomonas*, *Bacillus* and *Vibrio* species. Other than bacteria, piscidins have activity against fungi (Niu *et al.* 2013), parasites (Colorni *et al.* 2008), and viruses (Wang *et al.* 2010). Piscidins have immunomodulatory capacity to strengthen the innate immune system by modulating certain immune system-related genes (Masso-Silva and Diamond 2014).

Hepcidins are cysteine-rich peptides with wide availability among vertebrates including mammals, reptiles, amphibians and fish. The first hepcidin in fish was identified and isolated from hybrid striped bass (Shike *et al.* 2002) and since then hepcidins have been identified in around 40 other teleosts. In fish two forms of hepcidins can be identified: HAMP1 and HAMP2. HAMP1 is the most abundant member and present in both actinopterygian and non-actinopterygian fish but HAMP2 has been found only in actinopterygian fish. Fish hepcidins are potent antimicrobials against a wide variety of bacterial fish pathogens such as *Streptococcus iniae*, *Yersinia* and *Pseudomonas*, even at low micromole level (Masso-Silva and Diamond 2014). These are also effective against




FIGURE 2. Biological roles played by different AMPs. (Adapted from Diamond *et al.* 2009.)

different fungi and viruses.

Defensins is the group of cysteine-rich, cationic antimicrobial peptides found in plants, fungi, invertebrates and vertebrates. Although defensins are of three different types (α -, β -, and θ -defensins) in mammals, the defensins in fish belong to the β -defensin group. β -defensins in fish were initially identified from zebrafish, fugu and puffer fish (Zou *et al.* 2007). Since

then many defensins have been identified in many other marine and freshwater fish species. Fish β -defensins are mostly abundant in skin, head kidney and spleen. The antimicrobial activity of this AMP was established against pathogenic bacteria of fish such as *A. hydrophila*, *Y. ruckeri*, *V. anguillarum* and *E. tarda* and different fish-specific viruses including Singapore grouper iridovirus (SGIV), nodavirus and viral haemorrhagic septicaemia virus (VHSV).

Cathelicidins is an unusual group of AMPs with varying structure and size and defined by a conserved domain called 'cathelin.' The first member of this group in fish was identified in Atlantic hagfish *Myxine glutinosa* (Uzzell *et al.* 2003). The fish cathelicidins characterized by the cathelin domain from different fish species can be subdivided into two groups: the linear peptides and those that exhibit a characteristic disulphide bond (Masso-Silva and Diamond 2014). Fish cathelicidins have antimicrobial activity against fish pathogens such as *Y. ruckeri*, *A. salmonicida*, *C. albicans* and *Saprolegnia parasitica*.

APPLICATION AS THERAPEUTICS

The therapeutic potential of AMPs was evaluated for many critical diseases in mammals and recently these were evaluated for their potential to treat microbial infections in aquaculture. Among piscidins, oral and injection administration or electro-transfer of epinecidin-1 leads to enhanced survival in zebrafish and grouper infected with *V. vulnificus* and *Streptococcus agalactiae* (Lin *et al.* 2009, Pan *et al.* 2011a, 2012a). Tilapia hepcidin reduces the bacterial count and thereby fish mortality during *V. vulnificus* infection (Pan *et al.* 2012b). Transgenic zebrafish bearing tilapia hepcidin have enhanced bacterial resistance to *V. vulnificus* and *S. agalactiae* (Pan *et al.* 2011b).

CONCLUSION

Large-scale microbial infection in aquaculture frequently leads to considerable economic losses, as there are very few approved drugs available to counteract such a problem. Harnessing the potential of fish AMPs will give us a novel natural antimicrobial drug to handle this crisis situation against antibiotic-resistant microbes. More pronounced efforts are needed in the future to convert this baseline information into a potent, multifaceted, cost-effective drug for aquaculture.

Notes

- Anutosh Paria, ICAR- National Bureau of Fish Genetic Resources, Lucknow - 226 002, India. anu.cife@gmail.com; anutosh.paria@icar.gov.in
- Vinay T.N., ICAR- Central Institute of Brackishwater Aquaculture, Chennai - 600 028, India.
- Sanjay K. Gupta and Biplab Sarkar, ICAR- Indian Institute of Agricultural Biotechnology, Ranchi - 834 010, India.
- Tanmoy Gon Choudhury, College of Fisheries, Central Agricultural University (Imphal), Tripura - 799 210, India.

References

- Aoki, T. 1988. Drug resistance plasmids from fish pathogens. *Microbiological Sciences* 5:219-223.
- Buchanan, J.T., T.C. Cheng, J.F. LaPeyre, R.K. Cooper and T.R. Tiersch. 2001. *In vivo* transfection of adult eastern oysters *Crassostrea virginica*. *Journal of the World Aquaculture Society* 32:286-299.
- Cole, A.M., P. Weis and G. Diamond. 1997. Isolation and characterization of pleurocidin, an antimicrobial peptide in the skin secretions of winter flounder. *Journal of Biological Chemistry* 272:12008-12013.
- Colomi, A., A. Ullal, G. Heinisch and E.J. Noga. 2008. Activity of the antimicrobial polypeptide piscidin 2 against fish ectoparasites. *Journal of Fish Diseases* 31:423-432.
- DeGrandis, S.A. and R.M.W. Stevenson. 1985. Antimicrobial susceptibility patterns and R plasmid mediated resistance in the fish pathogen *Yersinia ruckeri*. *Antimicrobial Agents and Chemotherapy* 27:938-942.
- Diamond, G., N. Beckloff, A. Weinberg and K.O. Kisich. 2009. The roles of antimicrobial peptides in innate host defense. *Current Pharmaceutical Design* 15:2377-2392.
- FAO (Food and Agriculture Organization of the United Nations). 2016. *The State of World Fisheries and Aquaculture 2016, contributing to food security and nutrition for all*. Rome, Italy.
- Fischbach, M.A. and C.T. Walsh. 2009. Antibiotics for emerging pathogens. *Science* 325:1089-1093.
- Kathleen, M.M., L. Samuel, C. Felecia, E.L. Reagan, A. Kasing, M. Lesley and S.C. Toh. 2016. Antibiotic resistance of diverse bacteria from aquaculture in Borneo. *International Journal of Microbiology* 2016 (2164761):1-9.
- Lai, Y. and R.L. Gallo. 2009. AMPed up immunity: how antimicrobial peptides have multiple roles in immune defense. *Trends in Immunology* 30:131-141.
- Lee, C.S. and P.J. O'Bryen. 2003. Biosecurity in Aquaculture Production Systems: Exclusion of Pathogens and Other Undesirables. The World Aquaculture Society, Baton Rouge, Louisiana, USA.
- Lin, S.B., T.W. Fan, J.L. Wu, C.F. Hui and J.Y. Chen. 2009. Immune response and inhibition of bacterial growth by electrotransfer of plasmid DNA containing the antimicrobial peptide, epinecidin-1, into zebrafish muscle. *Fish and Shellfish Immunology* 26:451-458.
- Masso-Silva, J.A. and G. Diamond. 2014. Antimicrobial peptides from fish. *Pharmaceuticals* 7: 265-310.
- Mookherjee, N. and R.E. Hancock. 2007. Cationic host defence peptides: innate immune regulatory peptides as a novel approach for treating infections. *Cellular and Molecular Life Sciences* 64:922-933.
- Niu, S.F., Y. Jin, X. Xu, Y. Qiao, Y. Wu, Y. Mao, Y.Q. Su and J. Wang. 2013. Characterization of a novel piscidin-like antimicrobial peptide from *Pseudosciaena crocea* and its immune response to *Cryptocaryon irritans*. *Fish and Shellfish Immunology* 35:513-524.
- Pan, C.Y., J.L. Wu, C.F. Hui, C.H. Lin and J.Y. Chen. 2011a. Insights into the antibacterial and immunomodulatory functions of the antimicrobial peptide, epinecidin-1, against *Vibrio vulnificus* infection in zebrafish. *Fish and Shellfish Immunology* 31:1019-1025.
- Pan, C.Y., K.C. Peng, C.H. Lin and J.Y. Chen. 2011b. Transgenic expression of tilapia hepcidin 1-5 and shrimp chelonianin in zebrafish and their resistance to bacterial pathogens. *Fish and Shellfish Immunology* 31:275-285.
- Pan, C.Y., T.C. Huang, Y.D. Wang, Y.C. Yeh, C.F. Hui and J.Y. Chen. 2012a. Oral administration of recombinant epinecidin-1 protected grouper (*Epinephelus coioides*) and zebrafish (*Danio rerio*) from *Vibrio vulnificus* infection and enhanced immune-related gene expressions. *Fish and Shellfish Immunology* 32:947-957.
- Pan, C.Y., S.C. Lee, V. Rajanbabu, C.H. Lin and J.Y. Chen. 2012b. Insights into the antibacterial and immunomodulatory functions of tilapia hepcidin (TH) 2-3 against *Vibrio vulnificus* infection in mice. *Developmental and Comparative Immunology* 36:166-173.
- Pilet, M.F., X. Dousset, R. Barre, G. Novel, M. Desmazeaud and J.C. Piard. 1995. Evidence for 2 bacteriocins produced by *Carnobacterium piscicola* and *Carnobacterium divergens* isolated from fish and active against *Listeria monocytogenes*. *Journal of Food Protection* 58:256-262.
- Rajanbabu, V. and J.Y. Chen. 2011. Applications of antimicrobial peptides from fish and perspectives for the future. *Peptides* 32:415-420.
- Reddy K.V.R., R.D. Yedery and C. Aranha. 2004. Antimicrobial peptides: premises and promises. *International Journal of Antimicrobial Agents* 24:536-547.
- Sarmasik, A., G. Warr and T.T. Chen. 2002. Production of transgenic medaka with increased resistance to bacterial pathogens. *Marine Biotechnology* 4:310-322.
- Shike, H., X. Lauth, M.E. Westerman, V.E. Ostland, J.M. Carlberg, J.C. van Olst, C. Shimizu, P. Bulet and J.C. Burns. 2002. Bass hepcidin is a novel antimicrobial peptide induced by bacterial challenge. *European Journal of Biochemistry* 269:2232-2237.
- Stentiford, G.D., D.M. Neil, E.J. Peeler, J.D. Shields, H.J. Small, T.W. Flegel, J.M. Vlaskovits, B. Jones, F. Morado, S. Moss, J. Lotz, L. Bartholomay, D.C. Behringer, C. Hauton and D.V. Lightner. 2012. Disease will limit future food supply from the global crustacean fishery and aquaculture sectors. *Journal of Invertebrate Pathology* 110:141-157.
- The Fish Site. 2010. Aquatic animal diseases and their economic impact. 08 May 2010. <https://thefishsite.com/articles/aquatic-animal-diseases-and-their-economic-impact>.
- Uzzell, T., E.D. Stolzenberg, A.E. Shinnar and M. Zasloff. 2003. Hagfish intestinal antimicrobial peptides are ancient cathelicidins. *Peptides* 24:1655-1667.
- Wang, Y.D., C.W. Kung and J.Y. Chen. 2010. Antiviral activity by fish antimicrobial peptides of epinecidin-1 and hepcidin 1-5 against nervous necrosis virus in medaka. *Peptides* 31:1026-1033.
- Zou, J., C. Mercier, A. Koussounadis and C. Secombes. 2007. Discovery of multiple beta defensin like homologues in teleost fish. *Molecular Immunology* 44:638-647.